Children's Dental Village

Mercedes Padilla, D.D.S. Clinton Lepetich, D.M.D, M.S.D. 7360 S. McClintock Drive, Tempe AZ 85283 480-838-6949

Important Instructions for Lumineers

Please follow these guidelines:

 Your child must be very careful when eating and chewing. We ask that your child avoid biting into hard, crunchy or sticky foods and to avoid using the teeth to tear food. Eating these foods can cause the composite build-up to break or completely come off. Examples of foods to avoid include:

• **Hard foods**: Nuts, carrots, apples, hard candies, pizza crust and

ice (some foods can be cut into smaller pieces

and chewed with the back teeth).

Crunchy foods: Hard taco shells, chips, corn nuts.

• **Sticky foods**: Taffy, caramels, gummy bears, licorice and gum.

• **Tearing foods**: Jerky, corn on the cob, pizza crust (cut into smaller pieces and chew with back teeth).

- 2. Breakage can occur as a result of some force being placed on the tooth such as a bump to the mouth or hard foods. Accidents involving bumps to the teeth, night-time teeth grinding and chewing on items such as toys, pens and pencils, can also cause breakage or loss. Please be careful as there will be a fee to replace or repair the Lumineer.
- 3. The use of a mouth guard is recommended while playing sports.
- 4. Brush the Lumineer with a soft toothbrush and floss as you would your natural teeth, concentrating on the gum-line. Regular check-ups and cleanings are recommended to help insure the Lumineer and underlying tooth structures are healthy.
- 5. Your child may have been given a local anesthetic for their dental treatment. Children do not always understand the sensation of being numb. They may chew or suck on their lip, tongue or cheek and may cause painful wounds in their mouth.
- 6. We recommend a soft diet until the numbness wears off. Soft food suggestions include soup broth, Jello, pudding, yogurt, applesauce, ice cream or fruit juice.
- 7. If you have any questions, please do not hesitate to phone our office.